

Original Article

# PROLONGED INTERVAL BETWEEN TWIN LOSS AND TERM DELIVERY: A RARE CASE REPORT AND REVIEW OF THE LITERATURE

<sup>1</sup>*Adaeze Chinelo Nwankwo and* <sup>2</sup>*Ugochukwu Emmanuel Okeke*

<sup>1</sup>Departments of<sup>1</sup>Obstetrics and Gynaecology Chukwuemeka Odumegwu, University Teaching Awka, Anambra State, Nigeria.

<sup>2</sup>Obstetrics and Gynaecology Life, International Hospital, Awka, Nnamdi Azikiwe University Teaching Hospital, Nnewi, Anambra State.

DOI: <https://doi.org/10.5281/zenodo.13880406>

**Abstract:** While a number of cases detailing delayed interval birth of the second twin have been published recently, most of the deliveries have been before 34 weeks and rarely beyond 36 weeks. We report a case of delayed interval delivery at term with a favourable outcome. The loss of first twin was at 14 weeks in a dichorionic diamniotic twin gestation in which the placenta was retained. Pregnancy was conceived via In Vitro Fertilization (IVF) cycle. The Triangular 3-Bite Technique cervical stitch application was utilized to secure the cervix after 2 days of use of salbutamol and dydrogesterone as tocolytics with amoxicillian clavulanic acid to prevent intrauterine infection. The woman had a biweekly check of clotting profile and pregnancy was uneventful till term and she was delivered. We noted that delayed interval delivery can be done after the delivery of the first twin with second twin carried to term to avoid challenges of fetal prematurity.

**Keywords:** Delayed Interval Delivery, Twin Gestation, In Vitro Fertilization (IVF), Cervical Stitch, Fetal Prematurity

## Introduction

Gestational age at delivery has been reported as a strong predictor of neonatal survival with prematurity carrying a significant risk of increased neonatal morbidity and mortality.<sup>1</sup> Twin pregnancies are associated with higher incidence of preterm deliveries and the delivery of the second twin is usually expected to follow soon after the delivery of the first twin. However, delayed delivery for the second baby following loss of first twin has been reported as an option of management with possible good outcome.<sup>2</sup> Cases of successful delayed delivery of the second twin following prolongation of pregnancy until term have been a rarity. We report a rare case of delayed

## **Original Article**

interval delivery after application of cervical cerclage with subsequent prolongation of the second twin's delivery until term, achieved after a spontaneous abortion of first twin at 15 weeks gestational age.

### **CASE REPORT**

A 40-year old woman with dichorionic diamniotic twin pregnancy conceived via IVF. She booked for antenatal care at a gestational age of 13 weeks and was scheduled for cervical cerclage at 14 weeks but defaulted. However, at 15 weeks 2 days gestational age, she presented on account of severe lower abdominal pressure. Vaginal speculum examination showed a 3cm dilated cervix with obvious bleeding from the external os and bulging membrane. 2 hours later, she expelled the leading twin with the placenta retained for which it was ligated with vicryl 0 close to the external os. The patient was commenced on salbutamol 4mg tds and dydrogesterone 10mg tds as tocolysis and amoxicillin clavulanic acid injection (antibiotics). The patient and relatives were counseled on the option and consequences of a delayed interval delivery after cervical cerclage which they accepted. She subsequently had a Triangular 3-Bites Cervical Stitch, <sup>3</sup> applied on the cervix after 2 days with the placenta retained inside the uterus. She was maintained on salbutamol infusions for tocolysis, augmentin was continued for the next 24 hours to prevent intrauterine infections and she was closely monitored for 24 hours and discharged home on tabs augmentin and metronidazole for 1 week, hematinics and oral tocolytics. She was counseled on the need for regular antenatal care visit at two weeks interval with ultrasound follow up and evaluation of clotting profile.

She was regular with her visits and was closely monitored. She had her clotting time/prothrombin time, packed cell volume and urinalysis checked at every visit including ultrasound and vaginal swab culture and sensitivity done on monthly basis. She was placed on low dose aspirin, salbutamol tablet and hematinics till delivery. She had no clinical or laboratory signs of infections at any point.

Pregnancy remained uneventful and she had her baby via an elective cesarean section at a gestational age of 37 weeks with good outcome, weight was 3kg. The placenta of the dead twin was noted to be small, old and necrotic whereas that of the viable twin was normal. The cervical cerclage was removed intra-operative after delivery of the baby. Baby was transferred to mother in good condition.

Original Article

Placenta of the first twin noted to be small and atrophic at delivery



DISCUSSION

Twin pregnancies have been established to be associated with higher risk of preterm delivery than singletons and this carries significantly higher risk of morbidity and mortality.<sup>1</sup> The delivery of the second twin is routinely followed by the delivery of the second twin within a short interval either vaginally or by cesarean section. Delayed delivery after successful prolongation of gestational period with attendant increase in fetal weight in preterm cases for the second twin, significantly improve survival.<sup>2</sup>

Increased discussions and measures have been employed in recent years, in an attempt to prolong the pregnancy for the second twin after the first twin's delivery. A large number of cases with good results have been published. van Eyck et al,<sup>4</sup> in a prospective study, had obstetric follow-up of 50 cases of multiple pregnancies and reported that the delayed birth of the second twin is associated with reduced perinatal complications if the birth of the first twin happens between weeks 20 and 29. In a study of 35 patients, Fayad et al,<sup>5</sup> reported a mean interval of 47 days. A median duration of 6 days in a population of 200 pregnancies was reported by Benito Vielba et al.<sup>2</sup> Aydin and Celiloglu et al,<sup>6</sup> reported a case of delivery of an infant weighing 3639g delivered via cesarean section at 36 weeks' gestation with uneventful neonatal status after the first twin was delivered at 21weeks. Reports of varying intervals of delayed interval delivery have been documented in literature but that beyond 36weeks seem relatively scarce.

Varying degrees of complications may arise from delayed interval delivery including chorioamnionitis, coagulation disorder from retained placenta, abruption placenta, preterm labour among others but there is no established management guideline or protocol.<sup>5</sup> Chorioamnionitis following ascending infection from the ligated cord of first twin appears to be the earliest and commonest complication.<sup>5</sup> Its prevention appears to be the key to successful outcome. Empiric broad spectrum antibiotics should be initiated immediately as part of patient follow up. Periodic cervical and vaginal culture should be sampled and antibiotics administration based on the antibiogram instituted until negative culture result is obtained as was done in our patient.

## Original Article

In delayed interval delivery, one of the most controversial issues in management is the application of cervical cerclage. While some authors are averse to its use,<sup>2,4,7</sup> others found it important for a favourable outcome.<sup>3,5,6</sup> Generally, the application of cervical cerclage in twin and higher order pregnancies have remained controversial.<sup>8</sup> Its use have been recommended based on patient selection particularly when history based risk factors or ultrasound findings of cervical weakness are obtained<sup>9,10</sup> Our case suggests that patients need for cervical cerclage should be always individualized and patients' wish sort for. The couple gave consent for emergency cervical cerclage which helped to obtain a good outcome.

We simultaneously used two tocolytics in our patient initially with prolonged use of salbutamol only up till 34 weeks without adverse effect such as fetal tachycardia on the fetus as reported in literature. In our opinion, further studies are needed to gain knowledge in this regard.

We believe delayed interval delivery can be done after the delivery of the first twin with second twin carried to term to avoid challenges of fetal prematurity and its management especially in our environment where such services are not easily available.

## References

- Gladstone M, Oliver C, Van den Broek N. Survival, morbidity, growth and developmental delay for babies born preterm in low- and middle-income countries - a systematic review of outcomes measured. PLoS One. 2015 Mar 20;10(3):e0120566. doi: 10.1371/journal.pone.0120566. PMID: 25793703; PMCID: PMC4368095.
- Benito Vielba M, De Bonrosto Torralba C, Pallares Arnal V, Herrero Serrano R, Tejero Cabrejas EL, Campillos Maza JM. Delayed-interval delivery in twin pregnancies: report of three cases and literature review. J Matern Fetal Neonatal Med. 2019;32(2):351-355.
- Ikechebelu, JI1,2,3; Dim, CC4,5; Okpala, BC1,2,3; Eleje, GU1,2; Joe-Ikechebelu, NN3,6; Echezona, DM1; Nnoruka, MC3; Nwajiaku, LA3; Nwachukwu, CE7; Okam, PC1,3; Chigozie, AI3; Okpala, AN8; Igbodike, EP9. Comparison of Pregnancy Outcomes of Triangular 3-Bites and McDonald Techniques of Cervical Cerclage in Women with Cervical Insufficiency: A Pilot Study. Nigerian Journal of Clinical Practice 26(5):p 630635, May 2023. | DOI: 10.4103/njcp.njcp\_830\_22
- Arabin B, van Eyck J. Delayed-interval delivery in twin and triplet pregnancies: 17 years of experience in 1 perinatal center. Am J Obstet Gynecol. 2009;200(2):154.e1-154.e8.

**Original Article**

- Fayad S, Bongain A, Holhfeld P, et al. Delayed delivery of second twin: a multicentre study of 35 cases. *Eur J Obstet Gynecol Reprod Biol.* 2003;109(1):16-20.
- Aydin Y, Celiloglu M. Delayed interval delivery of a second twin after the preterm labor of the first one in twin pregnancies: delayed delivery in twin pregnancies. *Case Rep Obstet Gynecol.* 2012;2012:573824.
- Cozzolino M, Seravalli V, Masini G, Pasquini L, Di Tommaso M. Delayed-interval delivery in dichorionic twin pregnancies: A single-center experience. *Ochsner J.* 2015;15(3):248-250.
- Eleje GU, Eke AC, Ikechebelu JI, Ezebialu IU, Okam PC, Ilika CP. Cervical stitch (cerclage) in combination with other treatments for preventing spontaneous preterm birth in singleton pregnancies. *Cochrane Database Syst Rev.* 2020 Sep 24;9(9):CD012871. doi: 10.1002/14651858.CD012871.pub2. PMID: 32970845; PMCID: PMC809462
- Ikechebelu JI, Dim CC, Okpala BC, Eleje GU, Joe-Ikechebelu NN, Malachy DE, Nnoruka CM, Nwajiaku LA, Okam PC, Albert IC, Okpala AN, Igbodike EP. Comparison of Pregnancy Outcomes of History-Indicated and Ultrasound-Indicated Cervical Cerclage: A Retrospective Cohort Study. *Biomed Res Int.* 2023 Jan 9; 2023:8782854. doi: 10.1155/2023/8782854. PMID: 36654867; PMCID: PMC9842428.
- Frusca T, Zanardini C, Ghi T. Efficacy of ultrasound-indicated cerclage in twin pregnancies: is evidence-based medicine always the right choice? *Am J Obstet Gynecol.* 2016 Jan;214(1):131. doi: 10.1016/j.ajog.2015.08.057. Epub 2015 Sep 9. PMID: 26363482.